

Accessing the Gluon Polarization through A_{LL} in $\pi^\pm$ Production at PHENIX.

Astrid Morreale on behalf of the PHENIX collaboration

University of California, Riverside, CA, 92521 USA

Abstract. With proton-proton collisions at PHENIX a variety of direct channels are used to probe the proton substructure. Of the many channels available at PHENIX, charged pion measurements are expected to have sensitivity of Δg and thus help in the global analysis that will constrain it. We present a measurement of mid-rapidity charged pion production double longitudinal spin asymmetries (A_{LL}) at the p_T range of 5-10 GeV/c at collision energies of $\sqrt{s} = 200$ GeV.

Keywords: Double Longitudinal Asymmetry, Gluon, Δg , Proton Structure, Spin, Charged Pions, Pi Meson

PACS: 14.20.Dh, 25.40.Ep, 13.85.Ni, 13.88+e

INTRODUCTION

Measurements of charged pion asymmetries are an important process in the Δg global analysis that aims to disentangle all of the partonic contributions to the proton spin. In proton-proton collisions at PHENIX central rapidities, pion production proceed from quark-gluon and gluon-gluon initiated sub processes at the measured p_T range. This along with other pion meson properties such as zero spin and pseudo scalar under parity transformations, makes the pion an accessible channel whose importance may elucidate information on the gluon's contribution to the proton's spin.

π MESONS

Pions, being an isospin triplet, make combining asymmetry measurements from all three pion species for $5 < p_T < 12$ GeV/c particularly sensitivity to the sign of Δg as qg interactions dominate pion production in this p_T range. Preferential fragmentation of up quarks to positive pions and down quarks to negative pions leads to dominance of ug and dg contributions in the sum over flavors in a factorized pQCD calculation of pion production. In addition, the polarized parton distribution functions(pdf's) are well known to be $\Delta u > 0$ and $\Delta d < 0$ from polarized deep inelastic scattering (DIS) experiments. This dominance of u or d combined with the different signs on their polarized distributions translates into potentially measurable differences in the asymmetries for the different pion species that depend on the sign of Δg .

DETECTOR SETUP

The PHENIX detector at RHIC has fine-grained calorimetry 100 times finer than previous collider detectors, making particle identification excellent, the resolution of the electromagnetic calorimeter (EMCal) is $\delta\eta * \delta\phi = 0.01 * 0.01$. [1]. Triggering in the central arms allow us to select high p_T photons, electrons and charged pions. We select the charged pion signal by requiring a deposition of an energy cluster associated with a charged track in coincidence with the collision trigger(minimum bias trigger). Due to the hadronic response in the EMCal, less than $\frac{2}{3}$ of the charged pions trigger an event[2]. We furthermore identify charged pions by selecting particles that produced Čerenkov light in our ring imaging Čerenkov detector(RICH) but did not pass an electromagnetic shower-shape cut.

SIGNAL AND BACKGROUND

The measurement presented consists of 0.89 billion events analyzed within a p_T range of 5-10 GeV/c, corresponding to an analyzed sample of approximate 2.3 pb^{-1} , the average polarization was 47%. The primary source of potential background in this analysis comes from low energy electrons with misreconstructed momentum, and charged hadron tracks firing the RICH. Below 15 GeV/c the only particles that can produce light in the RICH(CO2 radiator) are: electrons(0.017 GeV/c), muons(3.5 GeV/c) and charged pions(4.7 GeV/c). We performed a series of detector cuts which include: EM-Cal shower-shape cut of prob < 0.2 designed to eliminate 80% of all electrons, a p_T -dependent energy cut to remove tracks with mis-reconstructed momentum and a cut of energy/momentum < 0.9 to remove the remaining electrons with nearly the correct reconstructed momentum. Muons are not considered a primary source of background for this analysis since primary muon to pion ratios have been previously measured to be less than 10^{-3} at PHENIX central rapidities. To estimate the remaining background fraction, the region below 4 GeV/c was fit to a power law. Extrapolating the fit under the signal region (solid line, turn on curve times powerlaw) the background is estimated to be < 5%.[Figure 1]

The A_{LL} and $\delta_{A_{LL}}$ formulae used in this analysis are:

$$A_{LL} = \frac{\sigma_{++} - \sigma_{+-}}{\sigma_{++} + \sigma_{+-}} = \frac{1}{|P_Y||P_B|} \frac{N_{++} - RN_{+-}}{N_{++} + RN_{+-}}, R = \frac{L_{++}}{L_{+-}} \quad (1)$$

$$\delta_{A_{LL}} = \frac{1}{|P_1||P_2|} \frac{2RN_{++}N_{+-}}{(N_{++} + RN_{+-})^2} \sqrt{\left(\frac{\Delta N_{++}}{N_{++}}\right)^2 + \left(\frac{\Delta N_{+-}}{N_{+-}}\right)^2} \quad (2)$$

Where $\sigma_{++}(\sigma_{+-})$ is the cross section with the beam in same(opposite) helicity configuration, $P_1(P_2)$ is defined as the polarization on separate beams and $N_{++}(N_{+-})$ is the particle yields with the beam in same(opposite) helicity configuration and L is the integrated luminosity.




FIGURE 1. p_T Spectrum of negative(left) and positive(right) charged pions fitted to a power law (dashed line) and functional form (solid line)




FIGURE 2. Measured negative(left) and positive(right) charged pion A_{LL} . Theory curves are GRSV max[3]($\Delta g = g$, red), GRSV std(black), GRSV min($\Delta g = -g$, blue).

RESULTS

The measurements for Run-05 double helicity asymmetries for positive and negative charged pions and the associated statistical errors are presented at figure 2. A scale uncertainty due to 20% uncertainty on each beam due to the polarization is not included. The statistical uncertainty in A_{LL} using Run-06 data currently in production is expected to be approximately 2.7 times smaller.

CONCLUSIONS

We presented a first measurement of A_{LL} at the p_T range 5-10 GeV. While the background in this measurement is significantly lower than π^0 [4], for example, current statistical uncertainties do not constrain Δg . The measurement, in particular the negative (although statistically limited) asymmetry observed in negative charged pions, begins

to hint to a possible problem in the fragmentation functions used in the parametrizations[Figure 2]. This measurement will be repeated with run-06 data as it becomes available. We expect that this analysis with the higher statistics and a cross-section measurement will be forthcoming and essential in pQCD interpretations and, ultimately, we expect to make a contribution to the global analysis that will determine Δg .

REFERENCES

1. K. Adcox *et al.*, *Nucl. Inst. Meth. A* **499**, 469 (2003).
2. J. Jiangyong, *High- p_T Charged Hadron Suppression in Au-Au Collisions at $\sqrt{s} = 200$ GeV*, Ph.D. thesis, Stony Brook University, New York, 11794 (2003).
3. M. Glück *et al.*, *Phys. Rev. D* **63**, 094005 (2001).
4. S. S. Adler *et al.*, *Phys. Rev. Lett.* **93**, 202002 (2004).